

AD-A247 576



Naval Health Research


2

**NORWEGIAN MILITARY FIELD EXERCISES IN THE ARCTIC:**

**ENDOCRINE AND METABOLIC RESPONSES**

*A. C. Hackney*

*J. A. Hodgdon*



*Report No. 91-9*

20030214264

92-06888



92 3 17 010

Approved for public release; distribution unlimited.

NAVAL HEALTH RESEARCH CENTER  
P.O. BOX 85122  
SAN DIEGO, CALIFORNIA 92186-5122

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND  
BETHESDA, MARYLAND



NORWEGIAN MILITARY FIELD EXERCISES IN THE ARCTIC:  
ENDOCRINE AND METABOLIC RESPONSES

A. C. Hackney, Ph.D.<sup>1</sup>

J. A. Hodgdon, Ph.D.<sup>2</sup>

<sup>1</sup>Exercise Physiology Laboratory  
Department of Exercise & Sport Science  
University of North Carolina  
Chapel Hill, North Carolina  
27599-8700

<sup>2</sup>Operational Performance Department  
Naval Health Research Center  
P. O. Box 85122  
San Diego, California  
92186-5122

Report No. 91-9, supported by the Naval Medical Research and Development Command, Department of the Navy under research work unit 63706N M0096.002-6003. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government. Approved for public release, distribution unlimited.

The authors wish to acknowledge the aid and support of the Norwegian Army, especially Company C, Brigade North. Also, to thank EM2 Brian Appleton, USN; HM3 Randy Howie, USN; HM2 Tina Jenkins, USN, Raymond Hilbert, and J. M. Shaw for their technical assistance.

### SUMMARY

This study examined the select endocrine and metabolic responses of Norwegian soldiers performing military field training while living in different shelter conditions in the arctic. A field based group (EXP, n = 17) lived in tents and a garrison group (CON, n = 16) lived in barracks for 10 days while performing similar daily training in cold weather. Cortisol, testosterone, thyroxine, glucose, triglycerides, and beta-hydroxybutyrate were measured on day 1, 5, and 10 of training.

The cortisol and testosterone findings suggest a moderate, but somewhat transient, stress response is associated with field living. The living conditions, however, did not seem to influence the metabolic responses to the physical activity. The observed changes in thyroxine, glucose, and triglycerides (i.e., reductions) suggest that an energy imbalance may accompany arctic military training.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## INTRODUCTION

Humans have extensive physiological defenses against heat, but very few and limited coping defenses against cold. In the cold they must rely heavily upon shelter and clothing to supplement and replace the innate physiological temperature regulation mechanisms of the body. That is, a strategy of behavioral temperature regulation is adopted. Many research studies of human responses to cold have been laboratory based where variables such as exposure duration, activity level, and clothing were rigidly controlled. Considerable extrapolation is required to apply these results to the field working conditions of cold regions because the natural thermal environments are complex and unstable, and individuals are free to modify their behavior to achieve comfort (i.e., adjust their clothing and activity for comfort) (1).

Military personnel of many nations periodically undergo physical training under an assortment of climatic conditions to improve their military readiness. Typically this military training is of a prolonged nature (i.e., several days). Cold weather exposure is a common environmental condition that is encountered with this training. Prolonged physical training of a military nature and cold exposure each are known to produce perturbations in the normal endocrine and metabolic function of the body (2,3,4,5). However, scanty information is available concerning the physiological consequences of prolonged field-based military training in a setting involving the cold environment encountered in the arctic region of the world. The intent of this study was to add to this limited knowledge base. Our purpose was twofold; 1) to measure select endocrine and metabolic changes to prolonged (10 day) military training in an arctic field setting involving cold exposure; and 2) to determine if shelter selection affected the physiological responses to the cold exposure.

## MATERIALS AND METHODS

Subjects. Our subjects were Norwegian military infantry soldiers who were all in good physical condition. All subjects gave informed consent prior to participation in the study. The subjects were divided into two groups. Group 1 (EXP; n = 17) lived in tents in the field; while group 2 (CON; n = 16) lived in a military garrison during the study. Throughout the study both groups carried out similar daily military operations (noted below), were equipped/clothed identically, and consumed the same food ration. Following completion of the study, three subjects were excluded from the CON group due to

methodological complications. Their physical characteristics (mean  $\pm$  SEM) were as follows: age; EXP =  $20.4 \pm 0.2$  yr, CON =  $20.5 \pm 0.3$  yr, height; EXP =  $182.3 \pm 1.7$  cm, CON =  $180.9 \pm 1.8$  cm; and weight EXP =  $79.7 \pm 2.0$  kg, CON =  $76.9 \pm 2.0$  kg.

Military Operations. The operations consisted of skiing and snowshoe marches, survival training, weapon training, and warfare maneuvers. The training took place in Skjold, Norway (approximately 300 km north of the arctic circle; elevation, 300 - 600 m) during late March, early April. The weather consisted of daily intermittent sunshine and overcast, with occasional blowing snow and temperatures of  $-20$  to  $+5^{\circ}$  C. Additionally, wind chill reductions of another 0 to  $-10^{\circ}$  C occurred frequently.

Blood Sampling & Analysis. Resting blood samples were taken on days 1, 5, and 10 of the study. The time of day for blood sampling was constant for each individual. The samples were obtained via venipuncture, and collected in an environmentally controlled field site ( $\sim 20 - 25^{\circ}$  C). Once collected, blood samples were kept cool ( $4^{\circ}$  C) until centrifugation ( $2000 \times g$ , 10 min). Separated sera was aliquoted and stored frozen at  $-50^{\circ}$  C until biochemical analysis. This analysis consisted of cortisol, testosterone, thyroxine, glucose, triglycerides, and beta-hydroxybutyrate (BOHB) determinations. All hormones were measured by radioimmunoassay procedures involving single anti-body, solid phase methodology (DPC Inc., Los Angeles, CA). All other blood measures were assayed with standard colorimetric procedures with reagents provided by Sigma Chemical Inc. (St. Louis, MO). Additionally, to assess the hydrational status of the subjects, serum osmolality was measured in all samples with a freeze point depression osmometer (Precision Systems Inc., Sudbury, MA). All biochemical analyses were performed in duplicate determinations.

The blood metabolite and hormones selected were felt to represent those physiological measures most likely to be affected by the experimental manipulation as based upon existing research findings (2,3,4,5).

Statistical Analysis. The data were analyzed with a mixed design (analysis of variance) with between group comparisons on the outcome measures and within person comparisons in the time domain. A Duncan multiple range test was used to determine individual mean differences and the significance level was set at  $p \leq 0.05$ . All reported values are means  $\pm$  SEM.

## RESULTS

The hormonal changes are depicted in Figures 1 - 3. Cortisol was unchanged in the CON group, but was increased significantly at day 10 from day 1 levels in the EXP group (see Fig. 1). Testosterone was also unchanged in the CON group, while in the EXP group a significant reduction was observed at day 5 from day 1 levels (see Fig. 2). However, by day 10 the testosterone levels of the EXP group were comparable to those found on day 1. No change was observed in the day 1 to 5 thyroxine levels. However, at day 10 a significant reduction in thyroxine occurred in both groups (see Fig. 3).

Table 1 reports the metabolic substrate findings. Glucose was significantly reduced at day 5 in both groups, but day 1 and day 10 levels were comparable, and within normal glycemia ranges. Triglyceride levels for both groups at days 5 and 10 were significantly lower than those found at day 1. Both groups showed a slight, but non-significant, elevation in BOHB by day 10. Also, no significant alterations were found in the serum osmolality for either group (see Table 1).

TABLE 1. Blood metabolic substrate responses  
of the experimental (EXP) and control (CON) groups  
during the 10 day military training in the arctic

Measure	Group	Day 1	Day 5	Day 10
<b>Glucose</b>				
(mg/dL)	EXP	69.5 ± 4.2	51.8 ± 3.2	77.0 ± 5.3
	CON	69.7 ± 5.4	49.6 ± 5.8	66.9 ± 10.1
<b>Triglycerides</b>				
(mg/dL)	EXP	199.9 ± 23.5	103.9 ± 13.7	91.7 ± 12.1
	CON	244.2 ± 28.1	168.4 ± 27.8	177.0 ± 25.9
<b>B-Hydroxybutyrate</b>				
(mg/dL)	EXP	8.9 ± 1.4	10.8 ± 2.2	13.2 ± 3.0
	CON	11.4 ± 2.0	13.8 ± 3.9	12.4 ± 3.2
<b>Osmolality</b>				
(mOsm/kg)	EXP	292.6 ± 1.5	289.5 ± 0.9	290.9 ± 1.0
	CON	290.8 ± 1.9	291.3 ± 1.4	293.7 ± 1.5

\* Means ± SEM

## DISCUSSION

The intent of our study was to measure the endocrine-metabolic changes to prolonged military field-based training in the arctic during cold exposure, and determine if shelter selections affected the physiological responses to the cold exposure.

The findings of elevated cortisol and reduced testosterone suggest a moderate, but somewhat transient, stress response is associated with living in the field shelter as opposed to living in the garrison (6,7). These findings are for the most part in agreement with previous work which has individually examined physiological responses to comparable military physical training and polar cold exposure (3,4,8). The observed thyroxine reduction may also be stress related; however, other possibilities are altered metabolic clearance or caloric deficiency (3,6,9). The last possibility seems more probable, since it is supported by the glucose and triglyceride findings. That is, the decreased glucose at day 5 may have been due to increased carbohydrate usage, a typical adaptation during the early stages of cold exposure (10). In turn the progressive reductions in triglyceride may represent a shift in the metabolism towards lipid substrate usage, with the reduced levels in the blood being due to increased tissue removal (5, 11). Although not significant, the change in BOHB suggests that an increased lipid metabolism may have been occurring in the subjects. Finally, the return of glucose to normal at day 10 may reflect a stabilization of carbohydrate balance as the body has shifted to lipid as a primary fuel source (11). Obviously this is speculation on our part, however, it is well established that military physical training imposes extreme caloric deficits, and this introduces metabolic alterations (2). Therefore, this hypothesis does present a feasible explanation of our data.

It is interesting that the living conditions did not seem to alter the metabolic substrate responses to the training in the arctic. Why the experimental treatment had an effect on certain of the hormones measured, but none of the metabolic substrates, is uncertain. Possibly if we had extended the duration of the experiment, the hormonal alterations noted would have precipitated detectable metabolism alterations. That is, there may have been insufficient time for the changes to become manifested.

We conclude that military physical training in arctic conditions does significantly alter the endocrine and metabolic status of soldiers. The

alterations noted may be driven by stress reactivity as well as an energy imbalance. Furthermore, the increased cold exposure encountered by living in the field during such training exercises shifts select aspects of the resting endocrine status; however, living conditions do not seem to impact blood metabolic substrate levels within the time frame of this study.



#### REFERENCES

1. Budd G.M. (1989). Ergonomic aspects of cold stress and cold adaptation. Scand. J. Work Environ. Health, 15(1), 15-26.
2. Opstad P.K., & Aakvaag A. (1983). The effect of sleep deprivation on the plasma levels of hormones during prolonged physical strain and calorie deficiency. Eur. J. Appl. Physiol., 51, 97-107.
3. Aakvaag A., Sand T., Opstad P.K., & Fonnum F. (1978). Hormonal changes in serum in young men during prolonged physical strain. Eur. J. Appl. Physiol., 39, 283-291.
4. Opstad P.K., & Aakvaag A. (1982). Decreased serum levels of oestradiol, testosterone and prolactin during prolonged physical strain and sleep deprivation, and the influence of a high calorie diet. Eur. J. Appl. Physiol., 49, 343-348.
5. Hurley B., & Haymes E. (1982). The effects of rest and exercise in the cold on substrate mobilization and utilization. Aviat. Space Environ. Med., 53, 1193-1197.
6. Mazzaferri, E.L. (ed.) (1985). Textbook of Endocrinology. New York: Medical Examination Publisher, 351-404.
7. Hackney, A.C. (1989). Endurance training and testosterone levels. Sports Med., 8, 117-127.
8. Reed H., Burman K., Shakir K., & O'Brian J. (1986). Alterations in the hypothalamic-pituitary-thyroid axis after prolonged residence in antarctica. Clin. Endocrinol., 25, 55-65.
9. Solter M., Brkic K., Petek M., Posavec L., & Seko M. (1989). Thyroid hormone economy in response to extreme cold exposure in healthy factory workers. J. Clin. Endocrinol. Metab., 68, 168-172.
10. Vallerand A.L., & Jacobs I. (1989). Rates of energy substrate utilization during human cold exposure. Eur. J. Appl. Physiol. 58, 873-878.
11. Shephard R.J. (1985). Adaptation to exercise in the cold. Sports Med., 2, 59-71.

#### FIGURE LEGEND

- FIGURE 1. The cortisol changes (mean  $\pm$  SEM) for the experimental (EXP) and control (CON) groups during the 10 days of arctic military training.
- FIGURE 2. The testosterone changes (mean  $\pm$  SEM) for the experimental (EXP) and control (CON) groups during the 10 days of arctic military training.
- FIGURE 3. The thyroxine changes (mean  $\pm$  SEM) for the experimental (EXP) and control (CON) groups during the 10 days of arctic military training.




Fig. 1




Fig. 2




Fig. 3

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</p>			
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE May 1991	3. REPORT TYPE AND DATE COVERED Interim Feb to Sept 1990	
4. TITLE AND SUBTITLE Norwegian Military Field Exercises in the Arctic: Endocrine and Metabolic Responses		5. FUNDING NUMBERS Program Element: 63706N Work Unit Number: M0096.002-6003	
6. AUTHOR(S) A. C. Hackney, J. A. Hodgdon		8. PERFORMING ORGANIZATION Report No. 91-9	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center P. O. Box 85122 San Diego, CA 92186-5122		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Medical Research and Development Command National Naval Medical Center Building 1, Tower 2 Bethesda, MD 20889-5044			
11. SUPPLEMENTARY NOTES Also: Proceedings - Nordic Conference on Cold, January 30 to February 2, 1991 Tromso, Norway			
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.		12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>This study examined the select endocrine and metabolic responses of Norwegian soldiers performing military field training while living in different shelter conditions in the arctic. A field based group (EXP, n = 17) lived in tents and a garrison group (CON, n = 16) lived in barracks for 10 days while performing similar daily training in cold weather. Cortisol, testosterone, thyroxine, glucose, triglycerides, and beta-hydroxybutyrate were measured on day 1, 5, and 10 of training.</p> <p>The cortisol and testosterone findings suggest a moderate, but somewhat transient, stress response is associated with field living. The living conditions, however, did not seem to influence the metabolic responses to the physical activity. The observed changes in thyroxine, glucose, and triglycerides (i.e., reductions) suggest that an energy imbalance may accompany arctic military training.</p>			
14. SUBJECT TERMS cold, military operations, endocrine response, metabolism		15. NUMBER OF PAGES	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE classified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited